

Exam 2 Nov. 10, 2005  
Math 25 Calculus I

SHOW ALL WORK  
Either circle your answers or place on answer line.

[16] 1.) If  $f'(x) = 3x^4 + 2 + 4x^{-1}$  and  $f(1) = 8$ , find  $f$

$$f(x) = \frac{3}{5}x^5 + 2x + 4\ln(x) + C \text{ for some } C.$$

$$f(1) = 8, f(1) = 8 = \frac{3}{5}(1)^5 + 2(1) + 4\ln(1) + C$$

$$8 = \frac{3}{5} + 2 + 0 + C$$

$$6 - \frac{3}{5} = \frac{30-3}{5} = \frac{27}{5} = C$$

$$\text{Answer 1.) } \underline{f(x) = \frac{3}{5}x^5 + 2x + 4\ln(x) + \frac{27}{5}}$$

[16] 2.)  $\lim_{x \rightarrow 0^+} [x^{x^3}] = \underline{1}$

$$\lim_{x \rightarrow 0^+} [x^{x^3}] = \lim_{x \rightarrow 0^+} [e^{\ln(x^{x^3})}] = \lim_{x \rightarrow 0^+} [e^{x^3 \ln(x)}]$$

$$\lim_{x \rightarrow 0^+} [x^3 \ln(x)] = \lim_{x \rightarrow 0^+} \left[ \frac{\ln(x)}{x^{-3}} \right] = \lim_{x \rightarrow 0^+} \left[ \frac{\frac{1}{x}}{-3x^{-4}} \right] = \lim_{x \rightarrow 0^+} \left[ \frac{x^3}{-3} \right] = 0$$

$$\text{Hence } \lim_{x \rightarrow 0^+} [e^{x^3 \ln(x)}] = e^{\lim_{x \rightarrow 0^+} [x^3 \ln(x)]} = e^0 = 1$$

[10] 3a.) Given  $x^2 + 2xy + y^3 - x - 3 = 0$ , then  $y' = \underline{\frac{1-2x-2y}{2x+3y^2}}$

$$\frac{d}{dx} (x^2 + 2xy + y^3 - x - 3) = \frac{d}{dx} (0),$$

$$2x + 2(y + xy') + 3y^2y' - 1 = 0,$$

$$2x + 2y + 2xy' + 3y^2y' - 1 = 0,$$

$$(2x + 3y^2)y' = 1 - 2x - 2y,$$

$$y' = \frac{1-2x-2y}{2x+3y^2},$$

[6] 3b.) Find the equation of the tangent line to the curve  $x^2 + 2xy + y^3 - x - 3 = 0$ , at the point  $(-2, 1)$ .

$$y' = \frac{1-2x-2y}{2x+3y^2},$$

$$\text{Hence at } (-2, 1), y' = \frac{1-2(-2)-2(1)}{2(-2)+3(1)^2} = \frac{1+4-2}{-4+3} = -3.$$

$$\frac{y-1}{x-(-2)} = -3, y = -3(x+2) + 1 = -3x - 6 + 1 = -3x - 5$$

$$\text{Answer 3b.) } \underline{y = -3x - 5}$$

[16] 4.) If  $g(3) = 4$  and  $g'(x) \leq 2$ , how large can  $g(8)$  be? 14

Since  $g'$  exists,  $g$  is differentiable and hence continuous.

By the MVT since  $g$  continuous on  $[3, 8]$  and  $g$  differentiable on  $(3, 8)$ , then there exists  $c \in [3, 8]$  such that

$$\frac{g(8) - g(3)}{8 - 3} = g'(c)$$

Hence  $\frac{g(8)-4}{5} = g'(c) \leq 2$ .

Thus  $g(8) \leq 2(5) + 4 = 14$ .

[16] 5.) A tank is in the form of an inverted cone having a height of 16m and a diameter at the top of 8 m. Water is flowing into the tank at the rate of  $2 \text{ m}^3/\text{min}$ . How fast is the water level rising when the water is 5m deep? (Volume of cone =  $\frac{1}{3}\pi r^2 h$ )

$$\frac{dV}{dt} = 2, \frac{dh}{dt} = ? \text{ when } h = 5.$$

NOTE:  $V$ ,  $h$ ,  $r$  change with respect to time. None of them are constant.

$$V = \frac{1}{3}\pi r^2 h, \quad \frac{r}{h} = \frac{4}{16}. \text{ Hence } r = \frac{1}{4}h.$$

$$V = \frac{1}{3}\pi\left(\frac{1}{4}h\right)^2 h = \frac{1}{3}\pi\left(\frac{h^3}{16}\right)$$

$$\frac{d}{dt}[V] = \frac{d}{dt}\left[\frac{1}{3}\pi\left(\frac{h^3}{16}\right)\right]$$

$$\frac{dV}{dt} = \frac{1}{3}\pi\left(\frac{3h^2}{16} \frac{dh}{dt}\right)$$

$$\frac{dV}{dt} = \pi\left(\frac{h^2}{16} \frac{dh}{dt}\right)$$

$$2 = \pi\left(\frac{5^2}{16} \frac{dh}{dt}\right). \text{ Hence } \frac{32}{25\pi} = \frac{dh}{dt}$$

Alternate answer:

$$V = \frac{1}{3}\pi r^2 h. \text{ NOTE: } V, h, r \text{ change with respect to time.}$$

$$\frac{d}{dt}[V] = \frac{d}{dt}\left[\frac{1}{3}\pi r^2 h\right]$$

$$\frac{dV}{dt} = \frac{1}{3}\pi\left[2r \frac{dr}{dt} h + r^2 \frac{dh}{dt}\right]$$

$$\text{When } h = 5: 2 = \frac{1}{3}\pi\left[2r \frac{dr}{dt}(5) + r^2 \frac{dh}{dt}\right]$$

$$\frac{r}{h} = \frac{4}{16}. \text{ Hence } r = \frac{1}{4}h. \text{ Thus } \frac{dr}{dt} = \frac{1}{4} \frac{dh}{dt} \text{ and when } h = 5, r = \frac{5}{4}.$$

$$\text{Thus, when } h = 5: 2 = \frac{1}{3}\pi\left[2\left(\frac{5}{4}\right)\left(\frac{1}{4}\right) \frac{dh}{dt}(5) + \left(\frac{5}{4}\right)^2 \frac{dh}{dt}\right] = \frac{25\pi}{16} \frac{dh}{dt}. \text{ Hence } \frac{32}{25\pi} = \frac{dh}{dt}$$

$$\text{Answer 5.) } \underline{\underline{\frac{dh}{dt} = \frac{32}{25\pi}}}$$

6.) Find the following for  $f(x) = \frac{x}{(x-1)^2}$  (if they exist; if they don't exist, state so). Use this information to graph  $f$ .

Note  $f'(x) = \frac{-x-1}{(x-1)^3}$  and  $f''(x) = \frac{2(x+2)}{(x-1)^4}$

[1] 6a.) critical numbers: -1

[1.5] 6b.) local maximum(s) occur at  $x =$  none

[1.5] 6c.) local minimum(s) occur at  $x =$  -1

[1.5] 6d.) The global maximum of  $f$  on the interval  $[0, 5]$  is none and occurs at  $x =$  none

[1.5] 6e.) The global minimum of  $f$  on the interval  $[0, 5]$  is 0 and occurs at  $x =$  0

[1.5] 6f.) Inflection point(s) occur at  $x =$  -2

[1] 6g.)  $f$  increasing on the intervals  $(-1, 1)$

[1] 6h.)  $f$  decreasing on the intervals  $(-\infty, -1) \cup (1, \infty)$

[1.5] 6i.)  $f$  is concave up on the intervals  $(-2, 1) \cup (1, \infty)$

[1.5] 6j.)  $f$  is concave down on the intervals  $(-\infty, -2)$

[1.5] 6k.) Equation(s) of vertical asymptote(s)  $x = 1$

[4] 6l.) Equation(s) of horizontal and/or slant asymptote(s)  $y = 0$

[4] 6m.) Graph  $f$

