

2.8: Approximating solution using

Method of Successive Approximation

(also called Picard's iteration method).

$$\text{IVP: } y' = f(t, y), y(t_0) = y_0.$$

Note: Can always translate IVP to move initial value to the origin and translate back after solving:

Hence for simplicity **in section 2.8**, we will assume initial value is at the origin: $y' = f(t, y), y(0) = 0$.

Thm 2.4.2: Suppose the functions

$z = f(t, y)$ and $z = \frac{\partial f}{\partial y}(t, y)$ are continuous on $(a, b) \times (c, d)$ and the point $(t_0, y_0) \in (a, b) \times (c, d)$,

then there exists an interval $(t_0 - h, t_0 + h) \subset (a, b)$ such that there exists a unique function $y = \phi(t)$ defined on $(t_0 - h, t_0 + h)$ that satisfies the following initial value problem:

$$y' = f(t, y), \quad y(t_0) = y_0.$$

Thm 2.8.1 is translated to origin version of Thm 2.4.2:

Thm 2.8.1: Suppose the functions

$z = f(t, y)$ and $z = \frac{\partial f}{\partial y}(t, y)$ are continuous for all t in $(-a, a) \times (-c, c)$,

then there exists an interval $(-h, h) \subset (-a, a)$ such that there exists a unique function $y = \phi(t)$ defined on $(-h, h)$ that satisfies the following initial value problem:

$$y' = f(t, y), \quad y(0) = 0.$$

Proof outline (note this is a constructive proof and thus the proof is very useful).

Given: $y' = f(t, y), y(0) = 0$ Eqn (*)
 $f, \partial f / \partial y$ continuous $\forall (t, y) \in (-a, a) \times (-b, b)$.

Then $y = \phi(t)$ is a solution to (*) iff

$$\phi'(t) = f(t, \phi(t)), \quad \phi(0) = 0 \text{ iff}$$

$$\int_0^t \phi'(s) ds = \int_0^t f(s, \phi(s)) ds, \quad \phi(0) = 0 \text{ iff}$$

$$\phi(t) = \phi(t) - \phi(0) = \int_0^t f(s, \phi(s)) ds$$

Thus $y = \phi(t)$ is a solution to (*)

$$\text{iff } \phi(t) = \int_0^t f(s, \phi(s)) ds$$

Construct ϕ using method of successive approximation – also called Picard’s iteration method.

Let $\phi_0(t) = 0$ (or the function of your choice)

Let $\phi_1(t) = \int_0^t f(s, \phi_0(s))ds$

Let $\phi_2(t) = \int_0^t f(s, \phi_1(s))ds$

⋮

Let $\phi_{n+1}(t) = \int_0^t f(s, \phi_n(s))ds$

Let $\phi(t) = \lim_{n \rightarrow \infty} \phi_n(t)$

To finish the proof, need to answer the following questions (see book or more advanced class):

- 1.) Does $\phi_n(t)$ exist for all n ?
- 2.) Does sequence ϕ_n converge?
- 3.) Is $\phi(t) = \lim_{n \rightarrow \infty} \phi_n(t)$ a solution to (*).
- 4.) Is the solution unique.

Example: $y' = t + 2y$. That is $f(t, y) = t + 2y$

Let $\phi_0(t) = 0$

$$\begin{aligned}\text{Let } \phi_1(t) &= \int_0^t f(s, 0)ds = \int_0^t (s + 2(0))ds \\ &= \int_0^t sds = \frac{s^2}{2} \Big|_0^t = \frac{t^2}{2}\end{aligned}$$

$$\begin{aligned}\text{Let } \phi_2(t) &= \int_0^t f(s, \phi_1(s))ds = \int_0^t f(s, \frac{s^2}{2})ds \\ &= \int_0^t (s + 2(\frac{s^2}{2}))ds = \frac{t^2}{2} + \frac{t^3}{3}\end{aligned}$$

$$\begin{aligned}\text{Let } \phi_3(t) &= \int_0^t f(s, \phi_2(s))ds = \int_0^t f(s, \frac{s^2}{2} + \frac{s^3}{3})ds \\ &= \int_0^t (s + 2(\frac{s^2}{2} + \frac{s^3}{3}))ds = \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{6}\end{aligned}$$

$$\begin{aligned}\text{Let } \phi_4(t) &= \int_0^t f(s, \phi_3(s))ds \\ &= \int_0^t f(s, \frac{s^2}{2} + \frac{s^3}{3} + \frac{s^4}{6})ds \\ &= \int_0^t (s + 2(\frac{s^2}{2} + \frac{s^3}{3} + \frac{s^4}{6}))ds \\ &= \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{6} + \frac{t^5}{15}\end{aligned}$$

⋮

Determine formula for ϕ_n :

Note patterns:

$$\int_0^t s ds = \frac{t^2}{2} =$$

$$\int_0^t \frac{s^2}{2} ds = \frac{t^3}{3 \cdot 2} =$$

$$\int_0^t \frac{s^3}{3 \cdot 2} ds = \frac{t^4}{4 \cdot 3 \cdot 2} =$$

$$\int_0^t \frac{s^4}{4 \cdot 3 \cdot 2} ds = \frac{t^5}{5 \cdot 4 \cdot 3 \cdot 2} =$$

Thus look for factorials.

$$\phi_0(t) = 0$$

$$\phi_1(t) = \frac{t^2}{2}$$

$$\phi_2(t) = \frac{t^2}{2} + \frac{t^3}{3}$$

$$\phi_3(t) = \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{6}$$

$$\phi_4(t) = \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{6} + \frac{t^5}{15} = \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{3 \cdot 2} + \frac{t^5}{5 \cdot 3}$$

Thus $\phi_n(t) =$

FYI (ie not on quizzes/exam):

$$\begin{aligned}\text{Defn: } \sum_{k=0}^{\infty} a_k x^k &= \lim_{n \rightarrow \infty} \sum_{k=0}^n a_k x^k \\ &= a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots\end{aligned}$$

Taylor's Theorem: If f is analytic at 0, then for small x (i.e., x near 0),

$$\begin{aligned}f(x) &= \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k \\ &= f(0) + f'(0)x + \frac{f''(0)}{2}x^2 + \frac{f'''(0)}{6}x^3 + \dots\end{aligned}$$

Example:

$$e^t = \sum_{k=0}^{\infty} \frac{t^k}{k!} \text{ and thus } e^{bt} = \sum_{k=0}^{\infty} \frac{b^k t^k}{k!} \text{ for } t \text{ near } 0.$$


$$\phi_n(t) = \sum_{k=2}^n \frac{2^{k-2}}{k!} t^k$$

$$\begin{aligned}\text{Thus } \phi(t) &= \lim_{n \rightarrow \infty} \phi_n(t) = \sum_{k=2}^{\infty} \frac{2^{k-2}}{k!} t^k = \frac{1}{4} \sum_{k=2}^{\infty} \frac{2^k}{k!} t^k \\ &= \frac{1}{4} (\quad - \quad - \quad)\end{aligned}$$

2.8: Approximating soln to IVP using seq of fns.

$$\phi_0(t) = 0, \quad \phi_1(t) = \frac{t^2}{2}, \quad \phi_2(t) = \frac{t^2}{2} + \frac{t^3}{3},$$

$$\phi_3(t) = \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{6}, \quad \phi_4(t) = \frac{t^2}{2} + \frac{t^3}{3} + \frac{t^4}{6} + \frac{t^5}{15}$$



2.7: Approximating soln to IVP using multiple tangent lines.

$$y(t) = \begin{cases} 0 & 0 \leq t \leq 0.1 \\ 0.1t - 0.01 & 0.1 \leq t \leq 0.2 \\ 0.22t - 0.034 & 0.2 \leq t \leq 0.3 \\ 0.364t - 0.0772 & 0.3 \leq t \leq 0.4 \\ 0.5328t - 0.14672 & 0.4 \leq t \leq 0.5 \end{cases}$$

