


[10] 1a.) Draw the direction field for the following differential equation:

$$y' = (y - 2)(y + 1)^2$$



[4] 1b.) On the direction field above, draw the solution to the above differential equation that satisfies the initial condition $y(1) = 0$.

[6] 1c.) Does the differential equation whose direction field is given above have any equilibrium solutions? If so, state whether they are stable, semi-stable or unstable.

Equilibrium solution = constant solution, $y = c$ and thus $y' = 0$

$$(y - 2)(y + 1)^2 = 0 \text{ implies } y = 2, -1$$

$y = 2$ is unstable, while $y = -1$ is semi-stable.

[15] 2.) Solve the initial value problem **for** y : $y' + \frac{3x}{y-4} = 0$, $y(1) = -2$.

$$\frac{dy}{dx} = -\frac{3x}{y-4}$$

$$\int (y-4)dy = \int -3xdx$$

$$\frac{y^2}{2} - 4y = -\frac{3}{2}x^2$$

$$y^2 - 8y = -3x^2 + C$$

$$y^2 - 8y + 3x^2 + C = 0$$

$$y = \frac{8 \pm \sqrt{64 - 4(3x^2 + C)}}{2} = 4 \pm \sqrt{16 - 3x^2 + C} = y$$

$$y(1) = -2: -2 = 4 \pm \sqrt{16 - 3(1)^2 + C} \text{ implies } -6 = -\sqrt{16 - 3 + C}$$

Note initial value determines sign of \pm . In this case, IVP only has a solution when we choose the negative sign. The the IVP in this case means $y = 4 - \sqrt{16 - 3x^2 + C}$ where we determine C below:

$$36 = 13 + C. \text{ Thus } C = 36 - 13 = 23 \text{ and } y = 4 - \sqrt{16 - 3x^2 + 23} = 4 - \sqrt{39 - 3x^2}$$

Answer: $y = 4 - \sqrt{39 - 3x^2}$

3.) Suppose $y' = y - t + 1$, $y(0) = 0$.

Let $\phi_0(t) = 0$ and define $\{\phi_n(t)\}$ by the method of successive approximation (i.e, Picards iteration method). Determine the following:

$$y' = f(t, y)$$

$$\phi_1(t) = \int_0^t f(s, \phi_0(s)) ds = \int_0^t f(s, 0) ds = \int_0^t (0 - s + 1) ds =$$

$$\left(-\frac{s^2}{2} + s\right)\Big|_0^t = -\frac{t^2}{2} + t - 0$$

$$[3] \quad 3a) \quad \phi_1(t) = \underline{-\frac{t^2}{2} + t}$$

$$\phi_2(t) = \int_0^t f(s, \phi_1(s)) ds = \int_0^t f(s, -\frac{s^2}{2} + s) ds = \int_0^t \left(-\frac{s^2}{2} + s - s + 1\right) ds$$

$$= \int_0^t \left(-\frac{s^2}{2} + 1\right) ds = \left(-\frac{s^3}{6} + s\right)\Big|_0^t = -\frac{t^3}{6} + t - 0$$

$$[3] \quad 3b) \quad \phi_2(t) = \underline{-\frac{t^3}{6} + t}$$

$$\phi_3(t) = \int_0^t f(s, \phi_2(s)) ds = \int_0^t f(s, -\frac{s^3}{6} + s) ds = \int_0^t \left(-\frac{s^3}{6} + s - s + 1\right) ds$$

$$= \int_0^t \left(-\frac{s^3}{6} + 1\right) ds = \left(-\frac{s^4}{24} + s\right)\Big|_0^t = -\frac{t^4}{24} + t - 0$$

$$[3] \quad 3c) \quad \phi_3(t) = \underline{-\frac{t^4}{24} + t}$$

$$[4] \quad 3d) \quad \phi_n(t) = \underline{-\frac{t^{n+1}}{(n+1)!} + t}$$

$$[3] \quad 3e) \quad \lim_{n \rightarrow \infty} \phi_n(t) = \underline{t}$$

[2] 3f) Is $\phi(t) = \lim_{n \rightarrow \infty} \phi_n(t)$ a solution to $y' = y - t + 1$, $y(0) = 0$? yes

[2] 3g) Is $\phi(t) = \lim_{n \rightarrow \infty} \phi_n(t)$ the unique solution to $y' = y - t + 1$, $y(0) = 0$? yes

[15] 4a.) Solve $y'' - 8y' + 16y = 0$

Educated guess: $y = e^{rt}$. Then $y' = re^{rt}$ and $y'' = r^2e^{rt}$

Plugging in the guess into our equation:

$$r^2e^{rt} - 8re^{rt} + 16e^{rt} = 0$$

Since $e^{rt} > 0$, we can divide both sides of the above equation by re^{rt} without losing any solutions:

$$r^2 - 8r + 16 = 0 \text{ implies } (r - 4)^2 = 0 \text{ and thus } r = 4.$$

Thus $y = e^{4t}$ is a solution. We can check by plugging in (as we did in class for a different example) that $y = te^{4t}$ is also a solution.

Sidenote: $\{e^{4t}, te^{4t}\}$ is a linear independent set and thus a basis for our solution. We can check linear independence by calculating the Wronskian.

$$\text{Answer: } \underline{y = c_1e^{4t} + c_2te^{4t}}$$

[15] 4b.) Solve $y'' - y' + 3y = 0$

Educated guess: $y = e^{rt}$. Then $y' = re^{rt}$ and $y'' = r^2e^{rt}$

Plugging in the guess into our equation:

$$r^2e^{rt} - re^{rt} + 3e^{rt} = 0$$

Since $e^{rt} > 0$, we can divide both sides of the above equation by re^{rt} without losing any solutions:

$$r^2 - r + 3 = 0 \text{ implies } r = \frac{1 \pm \sqrt{1-4(3)}}{2} = \frac{1 \pm \sqrt{11}i}{2} = \frac{1 \pm i\sqrt{11}}{2}.$$

$$\text{Answer: } \underline{y = c_1e^{\frac{1+i\sqrt{11}}{2}t} \cos\left(\frac{\sqrt{11}}{2}t\right) + c_2e^{\frac{1-i\sqrt{11}}{2}t} \sin\left(\frac{\sqrt{11}}{2}t\right)}$$

[15] 5.) Let $y = y_1(t)$ be a solution of $y' + p(t)y = 0$ and let $y = y_2(t)$ be a solution of $y' + p(t)y = g(t)$. Show that $y = y_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$.

Proof: Since $y = y_1(t)$ is a solution of $y' + p(t)y = 0$, we know that $y_1' + p(t)y_1 = 0$.

Since $y = y_2(t)$ is a solution of $y' + p(t)y = g(t)$, $y_2' + p(t)y_2 = g(t)$

Claim: $y = y_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$.

We will plug $y = y_1(t) + y_2(t)$ into the LHS to determine that the LHS = RHS:

$$\begin{aligned}(y_1(t) + y_2(t))' + p(t)(y_1(t) + y_2(t)) &= y_1'(t) + y_2'(t) + p(t)y_1(t) + p(t)y_2(t) \\ &= [y_1'(t) + p(t)y_1(t)] + [y_2'(t) + p(t)y_2(t)] = 0 + g(t) = g(t)\end{aligned}$$

Hence $y = y_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$.

Alternate proof: Since $y = y_1(t)$ is a solution of $y' + p(t)y = 0$, we know that

$$y_1' + p(t)y_1 = 0 \quad (1).$$

Since $y = y_2(t)$ is a solution of $y' + p(t)y = g(t)$.

$$y_2' + p(t)y_2 = g(t) \quad (2).$$

If we add equations (1) and (2), we obtain:

$$[y_1'(t) + p(t)y_1(t)] + [y_2'(t) + p(t)y_2(t)] = 0 + g(t)$$

Thus $y_1'(t) + y_2'(t) + p(t)y_1(t) + p(t)y_2(t) = g(t)$

and $(y_1(t) + y_2(t))' + p(t)(y_1(t) + y_2(t)) = g(t)$

Hence $y = y_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$.

Alternate proof:

Claim: $L(f) = f' + pf$ is a linear function where f and p are functions of t .

Proof of claim: Let a, b be constants and f, g be functions of t .

$$\begin{aligned}L(af + bg) &= (af + bg)' + p(af + bg) = af' + bg' + paf + pbg = af' + paf + bg' + pbg = \\ &= [a(f' + pf)] + [b(g' + pg)] = L(f) + L(g)\end{aligned}$$

We will now show that $y = y_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$:

Since $y = y_1(t)$ is a solution of $y' + p(t)y = 0$, $L(y_1) = 0$.

Since $y = y_2(t)$ is a solution of $y' + p(t)y = g(t)$, $L(y_2) = g(t)$

$L(y_1 + y_2) = L(y_1) + L(y_2) = 0 + g(t) = g(t)$. Thus $y = y_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$.

Note similar proofs would show that $y = cy_1(t) + y_2(t)$ is a solution of $y' + p(t)y = g(t)$ for any constant c .