

EXISTANCE

$\text{span}\{1, t, t^2\} =$ polynomials of degree ≤ 2

A polynomial $p(t)$ is in the span of $\{1, t, t^2\}$ if and only if there exists a solution for a, b, c to the equation

$$p(t) = a + bt + ct^2$$

Example 1: $2 + t^3$ is not in the span of $\{1, t, t^2\}$ since there does not exist a, b, c such that $2 + t^3 = a + bt + ct^2$

Example 2: $1 + 2t + 3t^2$ is in the span of $\{1, t, t^2\}$ since there exists a, b, c such that $1 + 2t + 3t^2 = a + bt + ct^2$

$a = 1, b = 2, c = 3$ is a solution.

UNIQUENESS

$\mathbf{b}_1, \dots, \mathbf{b}_n$ are linearly independent if

$$c_1\mathbf{b}_1 + c_2\mathbf{b}_2 + \dots + c_n\mathbf{b}_n = d_1\mathbf{b}_1 + d_2\mathbf{b}_2 + \dots + d_n\mathbf{b}_n$$

implies $c_1 = d_1, c_2 = d_2, \dots, c_n = d_n$.

or equivalently,

$\mathbf{b}_1, \dots, \mathbf{b}_n$ are linearly independent if

$c_1 \mathbf{b}_1 + c_2 \mathbf{b}_2 + \dots + c_n \mathbf{b}_n = 0$ implies $c_1 = c_2 = \dots c_n$.

Example 1: $\mathbf{b}_1 = (1, 0, 0)$, $\mathbf{b}_2 = (0, 1, 0)$, $\mathbf{b}_3 = (0, 0, 1)$. ■

$$(1, 2, 3) \neq (1, 2, 4).$$

If $(a, b, c) = (1, 2, 3)$ then $a = 1$, $b = 2$, $c = 3$.

Example 2: $\mathbf{b}_1 = 1$, $\mathbf{b}_2 = t$, $\mathbf{b}_3 = t^2$.

$$1 + 2t + 3t^2 \neq 1 + 2t + 4t^2.$$

If $a + bt + ct^2 = 1 + 2t + 3t^2$ then $a = 1$, $b = 2$, $c = 3$.

Application: Partial Fractions

$$\frac{4}{(x^2+1)(x-3)} = \frac{Ax+B}{x^2+1} + \frac{C}{x-3}$$

If you don't like denominators, get rid of them:

$$4 = (Ax + B)(x - 3) + C(x^2 + 1)$$

$$4 = Ax^2 + Bx - 3Ax - 3B + Cx^2 + C$$

$$4 = (A + C)x^2 + (B - 3A)x - 3B + C$$

I.e., $0x^2 + 0x + 4 = (A + C)x^2 + (B - 3A)x - 3B + C$

Thus $0 = A + C$, $0 = B - 3A$, $4 = -3B + C$.

$C = -A$, $B = 3A$, $4 = -3(3A) + -A \Rightarrow 4 = -10A$.

Hence $A = -\frac{2}{5}$, $B = 3(-\frac{2}{5}) = -\frac{6}{5}$, $C = \frac{2}{5}$.

$$\begin{aligned} \text{Thus, } \frac{4}{(x^2+1)(x-3)} &= \frac{-\frac{2}{5}x - \frac{6}{5}}{x^2+1} + \frac{\frac{2}{5}}{x-3} \\ &= \frac{-2x-6}{5(x^2+1)} + \frac{2}{5(x-3)} \end{aligned}$$

Alternatively, can plug in $x = 3$ to quickly find C and then solve for A and B . Can also use matrices to solve linear eqns.