

Charts for  $S^2$ :

Stereographic projection:

$$U_N = S^n - \{N = (1, 0, \dots, 0)\}.$$

$$\phi_N : U_N \rightarrow \mathbb{R}^n,$$

$$\phi_N(\mathbf{x}) = \frac{\mathbf{x}}{(1-x_0)}$$

$$U_S = S^n - \{S = (-1, 0, \dots, 0)\}.$$

$$\phi_S : U_S \rightarrow \mathbb{R}^n,$$

$$\phi_S(\mathbf{x}) = \frac{\mathbf{x}}{(1+x_0)}$$

Pre-atlas for  $S^n$ :  $\mathcal{A}_1 = \{(U_N, \phi_N), (U_S, \phi_S)\}$ ,

---

Projection:

$$U_{zp} = \{(x, y, z) \in S^2 \mid z > 0\},$$

$$\varphi_{zp} : U_{zp} \rightarrow \{(x, y) \mid x^2 + y^2 < 1\},$$

$$\varphi_{zp}(x, y, z) = (x, y)$$

$$U_{yp} = \{(x, y, z) \in S^2 \mid y > 0\},$$

$$\varphi_{yp} : U_{yp} \rightarrow \{(x, z) \mid x^2 + z^2 < 1\},$$

$$\varphi_{yp}(x, y, z) = (x, z)$$

$$U_{xp} = \{(x, y, z) \in S^2 \mid x > 0\},$$

$$\varphi_{xp} : U_{xp} \rightarrow \{(y, z) \mid y^2 + z^2 < 1\},$$

$$\varphi_{xp}(x, y, z) = (y, z)$$

$$U_{zn} = \{(x, y, z) \in S^2 \mid z < 0\},$$

$$\varphi_{zn} : U_{zn} \rightarrow \{(x, y) \mid x^2 + y^2 < 1\},$$

$$\varphi_{zn}(x, y, z) = (x, y)$$

$$U_{yn} = \{(x, y, z) \in S^2 \mid y < 0\},$$

$$\varphi_{yn} : U_{yn} \rightarrow \{(x, z) \mid x^2 + z^2 < 1\},$$

$$\varphi_{yn}(x, y, z) = (x, z)$$

$$U_{xn} = \{(x, y, z) \in S^2 \mid x < 0\},$$

$$\varphi_{xn} : U_{xn} \rightarrow \{(y, z) \mid y^2 + z^2 < 1\},$$

$$\varphi_{xn}(x, y, z) = (y, z)$$

Compatibility:

$$\varphi_{zp} \circ \varphi_{yn}^{-1} : \varphi_{yn}(U_{zp} \cap U_{yn}) \rightarrow \varphi_{zp}U_{zp} \cap U_{yn}).$$

$$U_{zp} \cap U_{yn} = \{(x, y, z) \in S^2 \mid z > 0 \text{ and } y < 0\}.$$

$$\varphi_{yn}(U_{zp} \cap U_{yn}) = \{(x, z) \mid z > 0 \text{ and } x^2 + z^2 < 1\}$$

$$\varphi_{zp} \circ \varphi_{yn}^{-1}(x, z) = \varphi_{zp}(x, -(1 - x^2 - z^2)^{1/2}, z) = (x, -(1 - x^2 - z^2)^{1/2}).$$

Pre-atlas for  $S^2$ :  $\mathcal{A}_2 = \{(U_{zp}, \varphi_{zp}), \dots, (U_{xn}, \varphi_{xn})\}$ ,

---

Polar coordinates

$$(x, y) = (r \cos \theta, r \sin \theta), r \geq 0, \theta \in [0, 2\pi),$$

$$(r, \theta) = (\sqrt{x^2 + y^2}, \tan^{-1}(y/x))$$

$$U_\gamma = S^1 - \{(1, \gamma)\},$$

$$\phi_{(1,\gamma)} : U_p \rightarrow (0, 2\pi)$$

$$\phi_{(1,\gamma)}(r, \theta) = \theta - \gamma \bmod 2\pi$$

Pre-atlas for  $S^1$ :  $\{(U_\gamma, \phi_{(1,\gamma)}) \mid \gamma \in S^1\}$

---

Spherical coordinates

$$(x, y, z) = (r \sin \phi \cos \theta, r \sin \phi \sin \theta, r \cos \phi), r \geq 0, \theta \in [0, 2\pi), \phi \in [0, \pi]$$

$$(r, \theta, \phi) = (\sqrt{x^2 + y^2 + z^2}, \tan^{-1}(y/x), \tan^{-1}(\sqrt{x^2 + y^2}/z)).$$

$$U_p = S^2 - \{p\},$$

$$\phi_p : U_p \rightarrow \phi_p(U)$$

$$\phi_p(r, \theta, \phi) = (\theta, \phi)$$


---

Cylindrical:  $(x, y, z) = (r \cos \theta, r \sin \theta, z), r \geq 0, \theta \in [0, 2\pi), z \in R$

$$(r, \theta, z) = (\sqrt{x^2 + y^2}, \tan^{-1}(y/x), z).$$

:

$$G(p, N) = \{[g] \mid g^{\text{smooth}} : U \rightarrow N, \text{ for some } U^{\text{open}} \text{ such that } p \in U \subset M\}$$

$$G(p) = G(p, \mathbf{R})$$

---

Let  $\alpha : I \rightarrow M$  where  $I = \text{an interval} \subset \mathbf{R}$ ,  $\alpha(0) = p$ . Note  $[\alpha] \in G[0, M]$

Directional derivative of  $[g]$  in direction  $[\alpha]$  =

$$D_\alpha g = \frac{d(g \circ \alpha)}{dt}|_{t=0} \in \mathbf{R}$$

---

$$T_p(M) = \{v : G(p) \rightarrow \mathbf{R} \mid v \text{ is linear and satisfies the Leibniz rule}\}$$

$v \in T_p(M)$  is called a *derivation*

---

Given a chart  $(U, \phi)$  at  $p$  where  $\phi(p) = \mathbf{0}$ , the *standard basis* for  $T_p(M) = \{v_1, \dots, v_m\}$ ,  
where  $v_i = D_{\alpha_i}$  and for some  $\epsilon > 0$ ,  $\alpha_i : (-\epsilon, \epsilon) \rightarrow M$ ,  $\alpha_i(t) = \phi^{-1}(0, \dots, t, \dots, 0)$

---

If  $v \in T_p(M)$ , then  $v = \sum_{i=1}^m a_i v_i$  where  $a_i = v([\pi_i \circ \phi])$

---

Suppose  $f^{\text{smooth}} : M \rightarrow N$ ,  $f(p) = q$ .

The *tangent (or differential) map*,

$$df_p : T_p M \rightarrow T_q N$$

$df_p(v) =$  the derivation which takes  $[g] \in G(q)$  to the real number  $v([g \circ f])$

I.e.,  $df_p$  takes the derivation  $(v : G(p) \rightarrow \mathbf{R}) \in T_p M$

to the derivation in  $T_q N$  which takes

the germ  $[g] \in G(q)$  to the real number  $v([g \circ f]).$