

Suppose $f : N \rightarrow M$ is smooth. Then $d_p f : T_p(N) \rightarrow T_p(M)$.

If df_p is 1 – 1, for all $p \in N$, then f is called an *immersion*.

I.e., f is an immersion iff f has rank n

If df_p is onto for all $p \in N$, then f is called a *submersion*.

I.e., f is an submersion iff f has rank m

Defn. Suppose $f : M \rightarrow N$ is smooth.

$p \in M$ is a *critical point* and $f(p)$ is a *critical value* if $\text{rank } df_p < n$.

If $p \in M$ is not a critical point, then it is a *regular point*.

If $q \in N$ is not a *critical value*, then it is a *regular value*.

Note: $q \in N$ is a regular value iff $f^{-1}(q) = \emptyset$ or $\forall p \in f^{-1}(q), df_p = n$.

Defn: If K is a submanifold of M , then K has the subspace topology.

Also, for all $p \in K$, there exists a chart (ϕ, U) for M such that $\phi(U) = \Pi_1^m(-\epsilon, \epsilon)$ and $\phi(U \cap K) = \Pi_1^k(-\epsilon, \epsilon) \times \Pi_{k+1}^m\{0\}$, then

Also, $(\phi|_{U \cap K}, U \cap K), \phi|_{U \cap K} : U \cap K \rightarrow \Pi_1^k(-\epsilon, \epsilon)$ is a chart for K .

The collection of such charts form a pre-atlas for K .

Thm 2.3.13: Let q be a regular value of $f: M \rightarrow N$. Then either $f^{-1}(q) = \emptyset$ or $f^{-1}(q)$ is an $(m - n)$ -submanifold of M .

Defn: Suppose $f: M \rightarrow N$ is a 1 – 1 immersion, and suppose $f: M \rightarrow f(M)$ is a homeomorphism, where $f(M) \subset N$ has the relative topology. Then f is an *embedding*, and $f(M)$ is an embedded submanifold.

Thm. $f : M \rightarrow N$ embedding implies $f(M)$ is a submanifold of N .