

6.3 Derangements

Suppose each person in a group of n friends brings a gift to a party. In how many ways can the n gifts be distributed so that each person receives one gift and no person receives their own gift.

Let the set of friends = $\{p_1, \dots, p_n\}$ where p_j = person j .

Let the set of gifts = $\{g_1, \dots, g_n\}$ where g_j = the gift brought by person j .

Suppose $f : \{p_1, \dots, p_n\} \rightarrow \{g_1, \dots, g_n\}$,

$f(p_k) = g_j$ iff person p_k receives give g_j , the gift brought by person j .

If each person receives one gift, then f is a bijection.

If no person receives their own gift. Then $f(p_j) \neq g_j$.

In simpler notation,

$f : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ such that $f(j) \neq j$

Recall:

a permutation on $\{1, \dots, n\}$ is a bijection $f : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$ ■

Ex: The permutation 1 2 3 4 5 corresponds to the identity function.

Ex: The permutation 1 3 2 corresponds to the function
 $f(1) = 1, f(2) = 3, f(3) = 2$

Defn: A *derangement* of $\{1, \dots, n\}$ is a permutation $i_1 i_2 \dots i_n$ such that $i_j \neq j$. I.e, j is not in the j th place.

In function notation: $f(j) = i_j$, then if $i_1 i_2 \dots i_n$ is a derangement, $f(j) \neq j$.

In yet other wording, recall a permutation corresponds to the placement of n non-attacking rooks on an $n \times n$ chessboard.

Ex: The permutation 1 3 2 corresponds to the following rook placement:

	●	
		●
●		

A derangement corresponds to non-attacking rook placement with forbidden positions along the diagonal (j, j) , for $j = 1, \dots, n$. ■

Ex: If rooks are placed on the following 3×3 chessboard in non-attacking position, then the rook placement corresponds to a derangement if no rook is placed in a spot marked with an X.

		X
	X	
X		

		X
	X	
X		

Thus the derangements of $\{1, 2, 3\}$ are 2 3 1 and 3 1 2.

Let $D_n =$ the number of derangements of $\{1, \dots, n\}$.

Thus $D_3 = 2$.

Thm 6.3.1: For $n \geq 1$, $D_n = n!(1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \dots + (-1)^n \frac{1}{n!})$

Pf: Use the inclusion and exclusion principle: If $A_i \subset S$,

$$\overline{\cup A_i} = |S| - \sum_{j=1}^n |A_j| + \sum_{i,j} |A_i \cap A_j| - \dots + (-1)^n |A_1 \cap A_2 \cap \dots \cap A_n|.$$

Choose S . What can we count which contains the set of derangements?

Let $S =$ the set of permutations of $\{1, \dots, n\}$. Then $|S| = n!$.

Choose A_j such that the set of derangements $= \overline{\cup A_j}$.

Let $A_j =$ set of permutations such that j is in the j th spot.

$|A_j| = (n - 1)!$ since there is only one choice for the j th spot (namely j), leaving $n - 1$ terms to permute in the remaining $n - 1$ places.

$|A_i \cap A_j| = (n - 2)!$ since there is only one choice for the i th spot (namely i) and only one choice for the j th spot (namely j), leaving $n - 2$ terms to permute in the remaining $n - 2$ places.

Similarly, $|A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}| = (n - k)!$.

Thus $D_n = n! - \sum_{j=1}^n (n - 1)! + \sum_{i,j} (n - 2)! - \dots + (-1)^n (n - n)!$

$$= \binom{n}{0} n! - \binom{n}{1} (n - 1)! + \binom{n}{2} (n - 2)! - \dots + \binom{n}{n} (-1)^n 0!$$

$$= n! - \frac{n!}{1!} + \frac{n!}{2!} + \dots + (-1)^n \frac{n!}{n!} = n!(1 - \frac{1}{1!} + \frac{1}{2!} + \dots + (-1)^n \frac{1}{n!})$$

Recall $\binom{n}{k} =$ number of ways to choose k A_i 's.

Sidenote: Finding the number of derangements is often called the hat check problem, because in the old days it was sometimes stated in the following terms: If n men check their hats, what is the probability that the hats are returned so that no one received their own hat.

Recall: If $E \subset S$, then the probability of $E = P(E) = \frac{|E|}{|S|}$

S = sample space, E = events.

Note: we assume each outcome is equally likely.

Suppose 4 customers at a restaurant order 4 meals. What is the probability that a waiter delivers these 4 orders to the 4 customers so that no customer receives what they ordered?

Answer: $\frac{D_4}{4!} = 1 - 1 + \frac{1}{2} - \frac{1}{6} = \frac{1}{3}$

The probability that a permutation of $\{1, \dots, n\}$ is a derangement $= \frac{D_n}{n!} = 1 - \frac{1}{1!} + \frac{1}{2!} + \dots + (-1)^n \frac{1}{n!}$

Recall Taylor's expansion from Calculus I,

$$f(x) = \sum_{j=0}^{\infty} (-1)^j \frac{f^{(n)}(a)}{j!}.$$

Thus $e^{-1} = \sum_{j=0}^{\infty} (-1)^j \frac{1}{j!}$.

Thus e^{-1} is a good approximation for the probability of a derangement for n (slightly) large.

Thus the probability of a derangement is about the same when $n = 5$ as it is for $n = 5000000000$.

We can derive a recursive formula for D_n (we will look at many

recursive formulas in chapter 7).

Lemma A: $D_n = (n - 1)(D_{n-2} + D_{n-1})$ for $n \geq 3$.

Note the above formula is a recursive formula as we can determine D_n by calculating D_k for $k < n$.

Note $D_1 = 0$, $D_2 = 1$ (as 2 1 is the only derangement of $\{1, 2\}$).

Thus $D_3 = 2(0 + 1) = 2$, $D_4 = 3(1 + 2) = 9$, $D_5 = 4(2 + 9) = 44$, etc.

Combinatorial proof of lemma A:

Let $\mathcal{D}_n =$ the set of derangements of $\{1, \dots, n\}$.

$D_n =$ the number of derangements of $\{1, \dots, n\} = |\mathcal{D}_n|$.

We need to show that D_n is a product of $n - 1$ and $D_{n-2} + D_{n-1}$. If we can partition \mathcal{D}_n into $n - 1$ subsets where each subset has $D_{n-2} + D_{n-1}$ elements, we can use the multiplication principle to show $D_n = (n - 1)(D_{n-2} + D_{n-1})$.

Let's focus on one of the positions of a derangement. The last (n th) position of our derangement can be anything except n . Thus there are $n - 1$ choices for the last (n th) position. Note the factor $n - 1$ appears in our formula.

Let $\mathcal{R}_k =$ the set of derangements of $\{1, \dots, n\}$ where k is in the n th position for $k = 1, \dots, n - 1$.

Then $\mathcal{D}_n = \cup_{j=0}^{n-1} \mathcal{R}_n$

Let $r_k = |\mathcal{R}_k|$ the number of derangements such that k is in the n th position.

Note that $r_1 = r_2 = \dots = r_{n-1}$ (while $r_n = 0$).

Then $D_n = r_1 + \dots + r_{n-1} = r_{n-1} + \dots + r_{n-1} = (n-1)r_{n-1}$.

Thus we have (hopefully) simplified our problem to showing that $D_{n-2} + D_{n-1} = r_{n-1} =$ the number of derangements such that $n-1$ is in the n th position.

We need to partition the permutations in \mathcal{R}_{n-1} into two sets, one with D_{n-2} elements and the other with D_{n-1} elements.

We can easily take care of D_{n-2} . The numbers $n-1$ and n do not appear in any derangement of $\{1, \dots, n-2\}$. In \mathcal{R}_{n-1} , $n-1$ appears in the last position. We can take a look at the derangements in \mathcal{R}_{n-1} , such that n appears in the $(n-1)$ st position. If we remove the n th and $(n-1)$ st entries, we obtain a derangement in \mathcal{D}_{n-2} .

Ex: for $n = 5$, $23154 \in \mathcal{R}_{n-1} \rightarrow 231 \in \mathcal{D}_{n-2}$.

Thus $D_{n-2} =$ the number of derangements of \mathcal{R}_{n-1} (such that $n-1$ is in the n th position and) n is in the $(n-1)$ st position.

We can now look at the remaining derangements in \mathcal{R}_{n-1} where n is not in the $(n-1)$ st position.

Let \mathcal{P}_n the set of derangement where $n-1$ is in the n th position and k is in the $(n-1)$ st position for some $k \neq n, n-1$ (I.e, $k \leq n-2$).

We would like to show that D_{n-1} = the number of derangements of $\{1, \dots, n-1\}$ such that $n-1$ is in the n th position and k is in the $(n-1)$ st position for some $k \leq n-2 = |\mathcal{P}_n|$.

Let \mathcal{D}_{n-1} = the set of derangements of $\{1, \dots, n-1\}$.

We would like to create a bijection from \mathcal{P}_n to \mathcal{D}_{n-1}

Note that the differences between \mathcal{P}_n and \mathcal{D}_{n-1} . A derangement in \mathcal{P}_n has n terms, while a derangement in \mathcal{D}_{n-1} has $n-1$ terms. Thus we need to remove a term to go from \mathcal{P}_n to \mathcal{D}_{n-1} .

If $i_1 i_2 \dots i_n \in \mathcal{P}_n$, then $i_n = n-1$ and $i_{n-1} = k$ for some $k \leq n-2$. Also $i_j = n$ for some j .

In \mathcal{D}_{n-1} , $i_{n-1} = k$ for some $k \leq n-2$ (by definition of derangement of $\{1, \dots, n-1\}$), so we have no problems with the $(n-1)$ st term.

However, we have the following differences between \mathcal{P}_n and \mathcal{D}_{n-1} :

$i_1 i_2 \dots i_n$ has n terms and

n appears somewhere in $i_1 i_2 \dots i_n$, and

$i_n = n-1$, so the placement of $n-1$ doesn't vary.

We can fix this by removing the n th term and replacing $i_j = n$ with $i_j = n-1$

Let $i_1 i_2 \dots i_n \in \mathcal{P}_n$. Then $i_n = n-1$ and $i_{n-1} = k$ for some $k \leq n-2$.

Create $a_1 a_2 \dots a_{n-1}$, a derangement of $\{1, \dots, n-1\}$ by

$$\text{let } a_l = \begin{cases} i_l & \text{if } i_l \neq n, 1 \leq l \leq n-1 \\ n-1 & \text{if } i_l = n \end{cases}$$

Ex: For $n = 5$, $25314 \in |\mathcal{P}_n| \rightarrow 2431 \in |\mathcal{D}_{n-1}|$.

This gives us a bijection between \mathcal{P}_n and \mathcal{D}_{n-1} . Thus $D_{n-1} = |\mathcal{P}_n|$.

Another (simpler) recurrence relation:

Lemma B: $D_n = nD_{n-1} + (-1)^n$ for $n \geq 2$

Proof by induction on n .

$n = 2$: $D_2 = 1$ (use definition or Thm 6.3.1)

$$2D_1 + (-1)^2 = 2(0) + 1 = 1.$$

Thus $D_n = nD_{n-1} + (-1)^n$ holds for $n = 2$.

Suppose $D_{k-1} = (k-1)D_{k-2} + (-1)^{k-1}$ for $k < n$

By lemma A, $D_k = (k-1)D_{k-2} + (k-1)D_{k-1}$

By the induction hypothesis, $D_{k-1} = (k-1)D_{k-2} + (-1)^{k-1}$.

$$\text{Thus } (k-1)D_{k-2} = D_{k-1} - (-1)^{k-1}$$

$$\text{Thus } D_k = D_{k-1} - (-1)^{k-1} + (k-1)D_{k-1} = kD_{k-1} + (-1)(-1)^{k-1} = kD_{k-1} + (-1)^k \blacksquare$$

6.4 Permutations with Forbidden Positions

Goal: To **derive** a more general formula for counting the number of permutations with arbitrary forbidden positions.

Recall in section 6.3, we looked at permutations with forbid-

den positions A derangement corresponds to non-attacking rook placement with forbidden positions along the diagonal (j, j) , for $j = 1, \dots, n$. In this section, we will cover arbitrary forbidden positions.

Let $X_j \subset \{1, \dots, n\}$ for $j = 1, \dots, n$.


Defn: $P(X_1, X_2, \dots, X_n)$ = the set of permutations $i_1 i_2 \dots i_n$ of $\{1, \dots, n\}$ such that $i_j \notin X_j$.

Defn: $p(X_1, X_2, \dots, X_n) = |P(X_1, X_2, \dots, X_n)|$

Ex: $P(X_1, X_2, \dots, X_n)$ corresponds to the set of derangements of $\{1, \dots, n\}$ if $X_j = \{j\}$. Thus $D_n = |P(\{1\}, \{2\}, \dots, \{n\})|$


Recall, we can visualize permutations with forbidden positions via $n \times n$ chessboards.

Ex: Derangements of $\{1, 2, 3\}$:
 $X_j = \{j\}$.



Non-derangement example:

$n = 4, X_i = \{j, j + 1\}, j = 1, 2, 3, X_4 = \emptyset$.



$$P(X_1, X_2, \dots, X_n) = P(\{1, 2\}, \{2, 3\}, \{3, 4\}, \emptyset) = \{3124, 3412, 3421, 4123\}.$$

$$p(X_1, X_2, \dots, X_n) = p(\{1, 2\}, \{2, 3\}, \{3, 4\}, \emptyset) \\ = |\{3124, 3412, 3421, 4123\}| = 4.$$

We can use the inclusion-exclusion principle to calculate $p(X_1, X_2, \dots, X_n)$ (although in many cases, the computation can be tediously long and beyond computer capabilities for large n).

Similar to the proof of Thm 6.3.1. By the inclusion-exclusion principle,

$$p(X_1, X_2, \dots, X_n) = |S| - \sum_{j=1}^n |A_j| + \sum_{i,j} |A_i \cap A_j| - \dots + (-1)^n |A_1 \cap A_2 \cap \dots \cap A_n|$$

where

Let $S =$ the set of permutations of $\{1, \dots, n\}$. Then $|S| = n!$.

Let $A_j =$ set of permutations $i_1 i_2 \dots i_n$ such that $i_j \in X_j$ (for a fixed j).

Note there are $|X_j|$ ways to place a rook in the j th position. There are $(n - 1)!$ ways to place the remaining $n - 1$ rooks so that the permutation belongs to A_j .

Thus $|A_j| = |X_j|(n - 1)!$.

$\sum_{j=1}^n |A_j| = \sum_{j=1}^n |X_j|(n - 1)! = (n - 1)! \sum_{j=1}^n |X_j| = r_1(n - 1)!$
where $r_1 = \sum_{j=1}^n |X_j|$.

Note $r_1 =$ number of ways to place 1 nonattacking rooks on an $n \times n$ chessboard so that the rook is in a forbidden position.

Let's now look at $A_j \cap A_k$. $i_1 i_2 \dots i_n \in A_j \cap A_k$, then $i_j \in X_j$ and $i_k \in X_k$. Thus there are $|X_j|$ ways to place a rook in the

j th position and $|X_k|$ ways to place a rook in the k th position. There are $(n - 2)!$ ways to place the remaining $n - 1$ rooks so that the permutation belongs to $A_j \cap A_k$.

Thus $|A_j \cap A_k| = |X_j||X_k|(n - 2)!$.

$\sum_{i,j} |A_i \cap A_j| = \sum_{i,j} |X_j||X_k|(n - 2)! = (n - 2)! \sum_{i,j} |X_j||X_k|$. Let $r_2 = \sum_{i,j} |X_j||X_k|$.

Note $r_2 =$ number of ways to place 2 nonattacking rooks on an $n \times n$ chessboard so that each of the 2 rooks is in a forbidden position.

Similarly, define $r_k =$ number of ways to place k nonattacking rooks on an $n \times n$ chessboard so that each of the k rooks is in a forbidden position.

Then $\sum |A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}| = r_k(k - 1)!$.

Thus we have proved:

Thm 6.4.1: $p(X_1, X_2, \dots, X_n) = n! - r_1(n - 1)! + r_2(n - 2)! - \dots + (-1)^n r_n$.

Note that if there are many forbidden positions, then r_k may be difficult to calculate and it may be easier to calculate $p(X_1, X_2, \dots, X_n)$ directly. If there are few forbidden positions, Thm 6.4.1 is the easier method to compute $p(X_1, X_2, \dots, X_n)$.

Examples:

Let $X = \{1, 2, 3\}$. $p(\{1, 2\}, \{1, 3\}, \{3\}) =$

Note in this case, it was easiest to count directly and not use

Thm 6.4.1.

Examples:

Let $X = \{1, 2, 3, 4, 5\}$. $p(\{1, 2\}, \{1, 3\}, \{3\}) =$

6.5 Another Forbidden Position Problem

Goal: To **derive** a formula for counting the number of permutations with relative forbidden positions.

Ex: Suppose children 1, 2, 3, 4, and 5 sit in a row in class. Children 1 and 2 cannot sit next to each other or they will cause trouble.

The order in which the children sit corresponds to a permutation of $\{1, 2, 3, 4, 5\}$. If 1 is in the i th spot, then 2 cannot be in the $i - 1$ st spot or the $i + 1$ th spot. Thus the pattern 21 or 12 cannot appear in our permutation. This is called a relative forbidden position as certain positions for the placement of 2 are forbidden, but these forbidden positions depend on the placement of 1.

We will focus on the relative forbidden position problem in which

Let $Q_n =$ the number of permutations of $\{1, 2, \dots, n\}$ in which none of the patterns 12, 23, 34, ..., $(n - 1)n$ occurs.

Thm 6.5.1 $Q_n = n! - \binom{n-1}{1} (n-1)! + \binom{n-1}{2} (n-2)! - \dots + \binom{n-1}{n-1} (-1)^{n-1} 1!$

Proof: Use inclusion-exclusion principle.

Let S = the set of permutations of $\{1, \dots, n\}$. Then $|S| = n!$.

Let A_j = set of permutations which contain the pattern $j(j+1)$.

Note: $|A_j| = (n - 1)!$

$|A_i \cap A_j| = (n - 2)!$

$|A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}| = (n - k)!$.