

Suppose f integrable

(Note f continuous implies f integrable).

If n equal subdivisions: $\Delta x = \frac{b-a}{n}$ and if we use right-hand endpoints: $x_i = a + i\Delta x = a + \frac{(b-a)i}{n}$

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(a + \frac{(b-a)i}{n}\right) \left(\frac{b-a}{n}\right)$$

Evaluate the limit by recognizing the sum as a Riemann sum for a function defined on $[0, 1]$

1.) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \sin\left(\frac{i}{n}\right) \frac{1}{n}$

2.) $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{i^5}{n^6}$

The Fundamental Theorem of Calculus: Suppose f continuous on $[a, b]$.

1.) If $G(x) = \int_a^x f(t)dt$, then $G'(x) = f(x)$.

2.) $\int_a^b f(t)dt = F(b) - F(a)$ where F is any antiderivative of f , that is $F' = f$.

The Fundamental Theorem of Calculus: Suppose f continuous on $[a, b]$.

1.) If $\frac{d}{dx} \left[\int_a^x f(t)dt \right] = f(x)$.

2.) $\int_a^b F'(t)dt = F(b) - F(a)$.

Examples:

1.) If $G_1(x) = \int_0^x t^2 dt$, then $G_1'(x) = \underline{\hspace{2cm}}$. ■

2.) If $G_2(x) = \int_5^x t^2 dt$, then $G_2'(x) = \underline{\hspace{2cm}}$. ■

3.) If $G_3(x) = \int_{-2}^x \sin(t^2) dt$, then $G_3'(x) = \underline{\hspace{2cm}}$. ■

4.) If $G_4(x) = \int_4^x \tan\left(\frac{t^3}{t+1}\right) dt$, then $G_4'(x) = \underline{\hspace{2cm}}$. ■

5.) If $G_5(x) = \int_1^x \sqrt{3t-5} dt$, then $G_5'(x) = \underline{\hspace{2cm}}$. ■

Examples:

1.) If $H_1(x) = \int_0^{x^3} t^2 dt$, then $H_1'(x) = \underline{\hspace{2cm}}$. ■

2.) If $H_2(x) = \int_5^{x^3} t^2 dt$, then $H_2'(x) = \underline{\hspace{2cm}}$. ■

3.) If $G(x) = \int_2^{\ln(x)} \frac{t}{t+1} dt$, then $G'(x) = \underline{\hspace{2cm}}$. ■

4.) If $G(x) = \int_3^{\frac{1}{x}} \sec(t) dt$, then $G'(x) = \underline{\hspace{2cm}}$. ■

The Fundamental Theorem of Calculus: Suppose f continuous on $[a, b]$.

1.) If $G(x) = \int_a^x f(t)dt$, then $G'(x) = f(x)$.

Proof

$$G'(x) = \lim_{h \rightarrow 0} \frac{G(x+h) - G(x)}{h},$$

$$= \lim_{h \rightarrow 0} \frac{\int_a^{x+h} f(t)dt - \int_a^x f(t)dt}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\int_x^{x+h} f(t)dt}{h}$$

$$\leq \lim_{h \rightarrow 0} \frac{\int_x^{x+h} M_h dt}{h}$$

where $M_h = \max\{f(t) \mid x \leq t \leq x+h\}$

(Note M_h exists by extreme value thm)

$$\leq \lim_{h \rightarrow 0} \frac{(M_h)(h)}{h}$$

$$\leq \lim_{h \rightarrow 0} M_h = f(x)$$

Similarly $G'(x) \geq f(x)$

(using $m_h = \min\{f(t) \mid x \leq t \leq x+h\}$)

The Fundamental Theorem of Calculus: Suppose f continuous on $[a, b]$.

2.) $\int_a^b f(t)dt = F(b) - F(a)$ where F is any antiderivative of f , that is $F' = f$.

Proof

Let $G(x) = \int_a^x f(t)dt$. Then $G'(x) = f(x)$ (ie, G is an antiderivative of f).

Let F be any antiderivative of f .

Then $F(x) = G(x) + C = \int_a^x f(t)dt + C$ for some constant C .

Thus $F(b) - F(a) = G(b) + C - [G(a) + C]$

$$= G(b) - G(a) = \int_a^b f(t)dt - \int_a^a f(t)dt = \int_a^b f(t)dt.$$

$$\begin{aligned} \int_0^\pi \cos^3(x)dx &= \int_0^\pi \cos^2(x)\cos(x)dx = \int_0^\pi (1 - \sin^2(x))\cos(x)dx \\ &= \int_0^0 (1 - u^2)du = 0 \end{aligned}$$

Let $u = \sin(x)$, $du = \cos(x)dx$,

when $x = 0$, $u = \sin(0) = 0$, when $x = \pi$, $u = \sin(\pi) = 0$

Shortcut method: Use symmetry.

For example:

If f is an odd function ($f(-x) = -f(x)$), then $\int_{-a}^a f(x)dx = 0$

If f is an even function, then $\int_{-a}^a f(x)dx = 2 \int_0^a f(x)dx$