Linear Functions

A function f is linear if $f(a\mathbf{x}+b\mathbf{y}) = af(\mathbf{x}) + bf(\mathbf{y})$

Or equivalently f is linear if
1.) $f(a\mathbf{x}) = af(\mathbf{x})$ and 2.) $f(\mathbf{x}+\mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$

Theorem: If f is linear, then $f(\mathbf{0}) = \mathbf{0}$

Proof: $f(\mathbf{0}) = f(0 \cdot \mathbf{0}) = 0 \cdot f(\mathbf{0}) = \mathbf{0}$

Example 1.) $f : R \to R$, $f(x) = 2x$

Proof:
$f(ax+by) = 2(ax+by) = 2ax+2by = af(x)+bf(y)$

Example 2.) $f : R^2 \to R^2$,
$f((x_1, x_2)) = (2x_1, x_1 + x_2)$

Proof: Let $\mathbf{x} = (x_1, x_2)$, $\mathbf{y} = (y_1, y_2)$

$a\mathbf{x}+b\mathbf{y} = a(x_1, x_2)+b(y_1, y_2) = (ax_1, ax_2)+(by_1, by_2) =$
$(ax_1 + by_1, ax_2 + by_2)$

$f(ax_1 + by_1, ax_2 + by_2)$

$= (2(ax_1 + by_1), ax_1 + by_1 + ax_2 + by_2)$

$= (2ax_1 + 2by_1, ax_1 + ax_2 + by_1 + by_2)$

$= (2ax_1, ax_1 + ax_2) + (2by_1, by_1 + by_2)$

$= a(2x_1, x_1 + x_2) + b(2y_1, y_1 + y_2)$

$= af((x_1, x_2)) + bf((y_1, y_2))$

Example 3.) D : set of all differential functions $\to$ set of all functions, $D(f) = f'$

Proof:
$D(af+bg) = (af+bg)' = af'+bg' = aD(f)+bD(g)$

Example 4.) Given a, b real numbers,
I : set of all integrable functions on [a, b] $\to R$,
$I(f) = \int_a^b f$

Proof: $I(sf + tg) = \int_a^b sf + tg = s\int_a^b f + t\int_a^b g =$
$sI(f) + tI(g)$

Example 5.) The inverse of a linear function is linear (when the inverse exists).

Suppose $f^{-1}(x) = c$, $f^{-1}(y) = d$.

Then $f(c) = x$ and $f(d) = y$ and
$f(ac + bd) = af(c) + bf(d) = ax + by$.

Hence $f^{-1}(ax + by) = ac + bd = af^{-1}(x) + bf^{-1}(y)$.

Example 6.) D : set of all twice differential functions $\rightarrow$ set of all functions, $L(f) = af'' + bf' + cf$

Proof:

$$\begin{aligned} L(sf + tg) &= a(sf + tg)'' + b(sf + tg)' + c(sf + tg) \\ &= saf'' + tag'' + sbf' + tbg' + scf + tcg \\ &= s(af'' + bf' + cf) + t(ag'' + bg' + cg) \\ &= sL(f) + tL(g) \end{aligned}$$

Consequence 1: If ψ_1, ψ_2 are solutions to $af'' + bf' + cf = 0$, then $3\psi_1 + 5\psi_2$ is also a solution to $af'' + bf' + cf = 0$,

Proof: Since ψ_1, ψ_2 are solutions to $af''+bf'+cf = 0$, $L(\psi_1) = 0$ and $L(\psi_2) = 0$.

$$\begin{aligned} \text{Hence } L(3\psi_1 + 5\psi_2) &= 3L(\psi_1) + 5L(\psi_2) \\ &= 3(0) + 5(0) = 0. \end{aligned}$$

Thus $3\psi_1+5\psi_2$ is also a solution to $af''+bf'+cf = 0$

Consequence 2:
If ψ_1 is a solution to $af'' + bf' + cf = h$
and ψ_2 is a solution to $af'' + bf' + cf = k$,
then $3\psi_1+5\psi_2$ is a solution to $af''+bf'+cf = 3h+5k$,

Since ψ_1 is a solution to $af''+bf'+cf = h$, $L(\psi_1) = h$.

Since ψ_2 is a solution to $af''+bf'+cf = k$, $L(\psi_2) = k$.

$$\begin{aligned} \text{Hence } L(3\psi_1 + 5\psi_2) &= 3L(\psi_1) + 5L(\psi_2) \\ &= 3h + 5k. \end{aligned}$$

Thus $3\psi_1 + 5\psi_2$ is also a solution to
$af'' + bf' + cf = 3h + 5k$

Calculus pre-requisites you must know.

Derivative = slope of tangent line = rate.

Integral = area between curve and x-axis (where area can be negative).

Integration by parts:

Derivative of a product: $(uv)' = uv' + vu'$

$$uv' = (uv)' - vu'$$
$$\int uv' = \int (uv)' - \int vu'$$
$$\int uv' = (uv) - \int vu'$$

Example: $\int e^{2x} sin(3x)$

Let $u = sin(3x)$, $dv = e^{2x}$

then $du = 3cos(3x)$, $v = \frac{1}{2}e^{2x}$

then $d^2u = -9sin(3x)$, $\int v = \frac{1}{4}e^{2x}$

$\int e^{2x} sin(3x) = \frac{1}{2} sin(3x)e^{2x} - \int \frac{3}{2} e^{2x} cos(3x)$

$= \frac{1}{2} sin(3x)e^{2x} - [\frac{3}{4} cos(3x)e^{2x} - \int \frac{-9}{4} sin(3x)e^{2x}$

$\int e^{2x} sin(3x) = \frac{1}{2} sin(3x)e^{2x} - \frac{3}{4} cos(3x)e^{2x} - \frac{9}{4} \int sin(3x)e^{2x}$

$\frac{13}{4} \int e^{2x} sin(3x) = \frac{1}{2} sin(3x)e^{2x} - \frac{3}{4} cos(3x)e^{2x}$

$\int e^{2x} sin(3x) = \frac{4}{13} [\frac{1}{2} sin(3x)e^{2x} - \frac{3}{4} cos(3x)e^{2x}]$

Exercise: Calculate $\int e^{x} cos(2x)$