

1.) Circle T for true and F for false.

[4] 1a.) If \vec{a} and \vec{b} are vectors in \mathbb{R}^3 , then $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$. T

[4] 1b.) The arc length s along the smooth curve with position vector $\vec{r}(t) = x(t)\vec{i} + y(t)\vec{j} + z(t)\vec{k}$ from $\vec{r}(a)$ to $\vec{r}(b)$ is, by definition

$$s = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2} dt \quad \text{T}$$

2.) Determine whether or not the four points $A(5, 2, -3)$, $B(6, 4, 0)$, $C(7, 5, 1)$, and $D(14, 14, 18)$ are coplanar. If not find the volume of the parallelepiped spanned by \vec{AB} , \vec{AC} , and \vec{AD} .

$$\vec{AB} = (6, 4, 0) - (5, 2, -3) = (1, 2, 3)$$

$$\vec{AC} = (7, 5, 1) - (5, 2, -3) = (2, 3, 4)$$

$$\vec{AD} = (14, 14, 18) - (5, 2, -3) = (9, 12, 21)$$

$$\begin{vmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 9 & 12 & 21 \end{vmatrix} = \begin{vmatrix} 1 & 2-2 & 3-3 \\ 2 & 3-4 & 4-6 \\ 9 & 12-18 & 21-27 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 0 \\ 2 & -1 & -2 \\ 9 & -6 & -6 \end{vmatrix} = \begin{vmatrix} -1 & -2 \\ -6 & -6 \end{vmatrix} = (-1)(-6) - (-6)(-2) = 6 - 12 = -6$$

Note most row/column operations change the determinant. Make sure you know which does and which does not.

Thus the volume of the parallelepiped spanned by \vec{AB} , \vec{AC} , and \vec{AD} is $|-6| = 6$.

1.) Circle T for true and F for false.

[4] 1a.) The partial derivative value $f_x(a, b)$ is the slope of a line tangent to a curve on which y is constant and which passes through the point $(a, b, f(a, b))$ on the surface $z = f(a, b)$. T

[4] 1b.) The graph of the function $f(x, y) = 2 - 3x + 4y$ is a plane. T

[12] 2.) Find the unit tangent and normal vectors to the curve $y = x^3$ at the point $(-1, -1)$.


$$y' = 3x^2$$

$y'(-1) = 3(-1)^2 = 3$. Thus slope of tangent line at $(-1, -1)$ is 3. Thus direction of tangent vector is (run, rise) = $(1, 3) = \left(\frac{dx}{dx}, \frac{dy}{dx}\right)$

Hence unit tangent vector = $\frac{(1,3)}{|(1,3)|} = \frac{(1,3)}{\sqrt{1^2+3^2}} = \frac{(1,3)}{\sqrt{10}} = \left(\frac{1}{\sqrt{10}}, \frac{3}{\sqrt{10}}\right)$

Normal vector is perpendicular to tangent vector.

Thus normal vector is either $\left(-\frac{3}{\sqrt{10}}, \frac{1}{\sqrt{10}}\right)$ or $\left(\frac{3}{\sqrt{10}}, -\frac{1}{\sqrt{10}}\right)$ as these are both unit vectors whose dot product with the tangent vector is 0. By picture below, the unit normal vector is $\left(\frac{3}{\sqrt{10}}, -\frac{1}{\sqrt{10}}\right)$



Thus unit tangent vector is $\left(\frac{1}{\sqrt{10}}, \frac{3}{\sqrt{10}}\right)$

unit normal vector is $\left(\frac{3}{\sqrt{10}}, -\frac{1}{\sqrt{10}}\right)$

1.) Circle T for true and F for false.

[4] 1a.) An equation for the plane through the three points $(2, 4, -3)$, $(3, 7, -1)$, $(4, 3, 0)$ is $11x + y - 7z = 46$ F

$$11(2) + 4 - 7(-3) = 26 + 21 = 47$$

$$11(3) + 7 - 7(-1) =$$

$$11(4) + 3 - 7(0) =$$

[4] 1b.) If the cost function $C(x, y)$ of a box with base of length x and height y is given by

$$C(x, y) = 0.1(xy + \frac{100}{y} + \frac{100}{x})$$

then C is an independent variable and x and y are dependent variables. F

[12] 2.) Find the arc length of the curve $x = \sin(2t)$, $y = \cos(2t)$, $z = 8t$ from $t = 0$ to $t = \pi$.

$$s = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2} dt = \int_0^\pi \sqrt{[2\cos(2t)]^2 + [-2\sin(2t)]^2 + [8]^2} dt$$

$$= \int_0^\pi \sqrt{4\cos^2(2t) + 4\sin^2(2t) + 64} dt = \int_0^\pi \sqrt{4 + 64} dt = \int_0^\pi \sqrt{68} dt = \int_0^\pi 4\sqrt{17} dt = 4\sqrt{17}t \Big|_0^\pi = 4\sqrt{17}\pi$$