

**Proposition 1.** *Let  $X$  be a CW complex.*

$$1. H_k(X^n, X^{n-1}) \cong \begin{cases} 0 & n \neq k \\ \mathbb{Z}^\ell & n = k \end{cases}$$

where  $\ell$  is the number of  $n$ -cells of  $X$  (potentially infinite).

2.  $H_k(X^n) = 0$  if  $k > n$ . In particular, if  $X$  is finite dimensional, then  $H_k(X) = 0$  for  $k > \dim X$ .

3. The inclusion  $i : X^n \hookrightarrow X$  induces

isomorphisms  $i_* : H_k(X^n) \rightarrow H_k(X)$  for  $k < n$ .

*Proof.* a.)  $H_k(X^n, X^{n-1}) \cong H_k(X^n/X^{n-1}) \cong \vee S^n$

b.)  $H_{k+1}(X^{n+1}, X^n) \rightarrow H_k(X^{n-1}) \rightarrow H_k(X^n) \rightarrow H_k(X^n, X^{n-1})$

and  $H_k(X^0) = 0$  for  $k > 0$  and for  $k \neq n$ ,  $H_k(X^{n-1}) \cong H_k(X^n)$ .

□

Given a CW complex  $X$ ,  $H_n^{CW}(X)$ , *cellular homology*, is the homology of the chain complex

$$\cdots \rightarrow H_{n+1}(X^{n+1}, X^n) \xrightarrow{d_{n+1}} H_n(X^n, X^{n-1}) \xrightarrow{d_n} H_{n-1}(X^{n-1}, X^{n-2}) \rightarrow \cdots$$

Note that  $H_n(X^n, X^{n-1}) = C_n^{CW}(X)$

$d_n(e_\alpha^n) = \sum_\beta d_{\alpha\beta} e_\beta^{n-1}$  where

$d_{\alpha\beta}$  is the degree of the map  $S_\alpha^{n-1} \rightarrow X^{n-1} \rightarrow S_\beta^{n-1}$

**Theorem 1.**  $H_{\bullet}^{CW}(X) \cong H_{\bullet}(X)$ .

The following computations follow.

$$H_{\bullet}(\Sigma_g) \cong \mathbb{Z}_{(0)} \oplus \mathbb{Z}_{(1)}^{2g} \oplus \mathbb{Z}_{(2)}$$

$$H_{\bullet}(N_g) \cong \mathbb{Z}_{(0)} \oplus (\mathbb{Z}^{g-1} \oplus \mathbb{Z}/2)_{(1)}$$

$$H_k(\mathbb{R}P^n) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0 \text{ and if } k = n \text{ is odd,} \\ \mathbb{Z}/2 & \text{if } k \text{ is odd and } 0 < k < n, \\ 0 & \text{otherwise.} \end{cases}$$

$$H_k(L_m(\ell_1, \dots, \ell_n)) \cong \begin{cases} \mathbb{Z} & \text{if } k = 0 \text{ or } 2n - 1, \\ \mathbb{Z}/m & \text{if } k \text{ is odd and } 0 < k < 2n - 1, \\ 0 & \text{otherwise.} \end{cases}$$

Defn: a *Moore space*, denoted by  $M(G, n)$ , is a simply connected CW complex  $X$  satisfying  $H_n(X) = G$  and  $\tilde{H}_i(X) \cong 0$  for  $i \neq n$ .

Example: If  $G$  is finitely generated, let  $X = (\vee S^n) \cup (\sqcup e_{\alpha}^{n+1})$ .

**Theorem 2.** *For finite CW complexes  $X$ , the Euler characteristic is*

$$\chi(X) = \sum_n (-1)^n \text{rank } H_n(X^n, X^{n-1}) = \sum_n (-1)^n \text{rank } H_n(X).$$

For example,

$$\chi(\Sigma_g) = 2 - 2g,$$

$$\chi(N_g) = 2 - g.$$